

Effect of Nipping and Varieties on Yield Attributes and Yield of Chickpea (*Cicer arietinum* L.)

K. Kushwaha¹, P.K. Tyagi², A.K. Shridhar³ and A. Tyagi⁴

^{1,3}PG Scholar (Agronomy), JNKVV, College of Agriculture, Tikamgarh (M.P.)

²Department of Agronomy, JNKVV, College of Agriculture, Tikamgarh (M.P.)

⁴PG Scholar (Agronomy), ITM, Gwalior (M.P.)

Corresponding author's email: pktyagi197071@yahoo.com

Abstract

Field experiment was conducted to study the effect of nipping practices and varieties on yield attributes and yield of chickpea during rabi 2020-21 at JNKVV, College of Agriculture, Tikamgarh (Madhya Pradesh). The experiment was conducted in split-plot design with three replications and comprised of three nipping practices viz., N1: no nipping, N2: nipping at 35 DAS and N3: nipping at 45 DAS as main plot treatments and four cultivars viz., V1: JG-12, V2: JG-36, V3: RVG 201 and V4: RVG 202 as sub-plot treatments. Results revealed that nipping of apical portion of chickpea at 45 DAS resulted into significantly more numbers of pods (103 plant^{-1}) and 100-seed weight (19.3 g) as compared to nipping at 35 DAS (89.6 plant^{-1} and 19.2 g, respectively) and the lowest yield attributes was observed with no nipping practices (82.4 plant^{-1} and 18.4 g, respectively). The significantly higher seed yield (2567 kg ha^{-1}), straw yield (4037 kg ha^{-1}), biological yield (6604 kg ha^{-1}) were also recorded with nipping practices at 45 DAS followed by nipping at 35 DAS (2331 kg ha^{-1} , 3581 kg ha^{-1} and 5912 , respectively) and the lowest (1945 kg ha^{-1} , 3377 kg ha^{-1} and 5322 kg ha^{-1} , respectively) with no nipping practices. Nipping in chickpea at 45 DAS increased the grain yield (kg ha^{-1}) by 32.0% and 10.1% over no nipping and nipping at 35 DAS, respectively. Among varieties, cv. JG 12 produced significantly more number of pods (102 plant^{-1}) followed by cvs. RVG 201 (94.9 plant^{-1}), JG 36 (89.9 plant^{-1}) and the lowest in cv. RVG 202 (80.0 plant^{-1}). The significantly higher seed yield (2506 kg ha^{-1}), straw yield (3862 kg ha^{-1}) and biological yield (6368 kg ha^{-1}) were also recorded in cv. JG 12 followed by cv. RVG 201 (2315 kg ha^{-1} , 3738 kg ha^{-1} and 6053 , respectively), JG 36 (2217 kg ha^{-1} , 3557 kg ha^{-1} and 5774 , respectively) and significantly the lowest in cv. RVG 202 (2086 kg ha^{-1} , 3502 kg ha^{-1} and 5588 kg ha^{-1} , respectively). Variety JG-12 increased the grain yield (kg ha^{-1}) by 8.2 %, 13.0% and 20.1% over RVG-201, JG-36 and RVG-202, respectively. The combination of nipping at 45 DAS with cv. JG-12 increased the grain yield (kg ha^{-1}) by 4.5% to 56.6% over rest of the combinations of treatments.

Key words: chickpea, nipping, yield, yield attributes, varieties

Introduction

Chickpea (*Cicer arietinum* L.) is the second-most main pulse crop after pigeon pea in the humankind for diet and other use. India is the leading producer of chickpea in the world giving out 65% area and 70% of total global production. India is the principle chickpea producing country in the world with a total production of 10.13 million tonnes and cultivated area of 9.44 million hectare with

an average yield of 1073 kg ha^{-1} . In Madhya Pradesh, it is cultivated over an area of 3.43 million hectare and total production of 4.61 million tonnes with an average yield of 1344 kg ha^{-1} . There are many factors responsible for the low yield of chickpea viz., utilization of conventional or low yielding varieties and adoption of poor management practices. Among the agronomic management practices, nipping

is one of the important practices for the enhancement of yield, yield attributes and economics of chickpea. Nipping means the removal of top portion (apical meristem) of a plant to induce branching on the plant at the remaining nodes. Nipping of crop at various growth stages led to enhance number of branches and number of pods that consecutively boost yield of chickpea. Nipping at 45 DAS in chickpea increased yield as well as controlled disease severity. Nipping practice in the research area has two fold advantages. On the one hand, nipping at prescribed growth stages could improve yield of the crop while on the opposite hand during time, the chickpea within the field is typically a shortage of fodder and poor farmers couldn't afford to purchase forage at distant locations, so chickpea may provide them a chance to fetch green fodder for his/her livestock.

Materials and methods

Field experiment was conducted at Agronomy Research Area, J.N.K.V.V., College of Agriculture, Tikamgarh, Madhya Pradesh (24° 43' N latitude and 78° 49' E longitude at an altitude of 358m above mean sea level) during *rabi* 2020-21. The experimental site is of sub-tropical climate characterized by hot dry summers and cool dry winter lies in the Bundelkhand Zone (Agro-climatic Zone-VIII). The soil of experimental field was medium to deep black and clayey loam in texture having pH 7.1, EC 0.12 dS m⁻¹, organic carbon 0.5%, available N 265 kg ha⁻¹, available P₂O₅ 26 kg ha⁻¹ and available K₂O 260 kg ha⁻¹. The average annual rainfall of this region is about 1000 mm, which is mostly received between June to September and a little rainfall of 90 mm is also obtained during October to

The crop growth and yield of chickpea varieties under nipping at various stages help to pick the foremost promising variety in term of yield potential. Grain, straw and biological yield of chickpea crop were significantly influenced by the different varieties and nipping. The yield diversity concerning different varieties may also be caused by variation in pod bearing ability and therefore the number of seeds per pod^[2]. There is highly significant difference in the interaction of chickpea varieties with nipping which indicates that different varieties of chickpea reacted in a different way to nipping^[7]. Keeping these facts in view, an experiment was conducted to study the effect of nipping on yield attributes and yield of chickpea varieties.

May. The average temperature ranges between 4.5°C to 45°C. The experiment was conducted in split-plot design with three replications and comprised of three nipping practices viz., N1; no nipping, N2: nipping at 35 DAS and N3: nipping at 45 DAS as main plot treatments and four cultivars viz., V1: JG-12, V2: JG-36, V3: RVG 201 and V4: RVG 202 as sub-plot treatments. The full recommended doses of nitrogen (20 Kg N ha⁻¹), phosphorus (40 Kg P₂O₅ ha⁻¹) and potassium (20 kg K₂O ha⁻¹) were applied at sowing. The chickpea crop was sown in lines 30 cm apart using a seed rate of 80 kg ha⁻¹. Nipping was done at 35 DAS and 45 DAS in the respective treatments. All other agronomic and plant protection measures were applied as per recommendations. Yield attributes were recorded from the five plants sample

collected at the time of harvest. The crop harvested from net plot area was threshed after 4-5 days of sun drying and the seed yield of net plot was then converted into kg ha^{-1} . Before threshing of the crop harvested from net plot, the sun dried whole plant samples were weighed and

Results and discussion

Effect of nipping practices on yield attributes and yield

Data pertaining to yield attributes and seed yield as affected by nipping practices has been given in Table 1 and Table 2, respectively. Results revealed that nipping practices in chickpea crop significantly influenced the yield attributing characters. Nipping of apical portion of chickpea at 45 DAS resulted into significantly more numbers of pods

then converted into kg ha^{-1} to obtain biological yield. Straw yield is obtained by subtracting seed yield (kg ha^{-1}) from biological yield (kg ha^{-1}). The results were analyzed statistically to draw suitable inference as per standard ANOVA technique^[5].

(103 plant^{-1}) and 100-seed weight (19.3 g) as compared to nipping at 35 DAS (89.6 plant^{-1} and 19.2 g, respectively) and the lowest yield attributes was observed with no nipping practices (82.4 plant^{-1} and 18.4 g, respectively). However, nipping practices at 35 and 45 DAS failed to significantly influence the number of seeds pod^{-1} .

Table 1 Effect of nipping practices and varieties on yield attributes of chickpea

Treatments	Number pods plant^{-1}	Number seeds pod^{-1}	Seed index (g)
Nipping			
N1 : No nipping	82.4	1.23	18.4
N2 : Nipping at 35 DAS	89.6	1.27	19.2
N3 : Nipping at 45 DAS	103	1.30	19.3
S.Em \pm	0.37	0.03	0.13
CD (P=0.05)	1.44	NS	0.52
Varieties			
V1 : JG 12	102	1.31	16.6
V2 : JG 36	89.9	1.25	16.9
V3 : RVG 201	94.9	1.27	21.7
V4: RVG 202	80.0	1.24	20.7
S.Em \pm	0.41	0.05	0.32
CD (P=0.05)	1.23	NS	0.95
Interaction (nipping x variety)			
N1V1	93.2	1.28	16.8
N1V2	78.6	1.34	16.7

N1V3	85.4	1.17	21.5
N1V4	72.5	1.14	18.6
N2V1	100	1.33	16.5
N2V2	90.7	1.20	16.5
N2V3	93.5	1.19	21.4
N2V4	74.1	1.35	22.5
N3V1	112	1.31	16.4
N3V2	101	1.22	17.5
N3V3	106	1.45	22.3
N3V4	93.3	1.22	20.9
S.Em \pm	0.72	0.09	0.55
CD (P=0.05)	2.13	NS	1.65

Table 2 Effect of nipping practices and varieties on yield of chickpea

Treatments	Seed yield (kg ha ⁻¹)	Straw yield (kg ha ⁻¹)	Biological yield (kg ha ⁻¹)	Harvest index (%)
Nipping				
N1 : No nipping	1945	3377	5322	36.5
N2 : Nipping at 35 DAS	2331	3581	5912	39.4
N3 : Nipping at 45 DAS	2567	4037	6604	38.9
S.Em \pm	18.4	52.8	53.8	0.41
CD (P=0.05)	72.3	207.5	211.2	1.62
Varieties				
V1 : JG 12	2506	3862	6368	39.3
V2 : JG 36	2217	3557	5774	38.3
V3 : RVG 201	2315	3738	6053	38.2
V4: RVG 202	2086	3502	5588	37.4
S.Em \pm	28.4	58.0	69.1	0.42
CD (P=0.05)	84.5	172.2	205.3	1.23
Interaction (nipping x variety)				
N1V1	2064	3386	5450	37.9
N1V2	1931	3367	5297	36.5
N1V3	2003	3397	5400	37.1
N1V4	1781	3358	5139	34.7
N2V1	2667	3861	6528	40.8

N2V2	2167	3522	5689	38.1
N2V3	2367	3767	6133	38.6
N2V4	2125	3172	5297	40.2
N3V1	2789	4339	7128	39.1
N3V2	2553	3783	6336	40.3
N3V3	2575	4050	6625	38.9
N3V4	2353	3975	6328	37.2
S.Em \pm	49.2	100.4	119.7	0.72
CD (P=0.05)	146.3	298.2	355.6	2.14

The increase in yield attributing parameters noticed with nipping practice was mainly due to accumulation of more photosynthates which were utilized for production of more number of productive secondary branches and more number of pods plant⁻¹ in chickpea. The nipping at 45 DAS might have efficiently altered the crop architecture by activating the dormant lateral secondary branches which ultimately increased the number to pods plant⁻¹ which leads to greater chance for development of source and sink relationship in chickpea and thereby would have facilitated the significant increase in the yield attributes of the chickpea. This has also been documented by other worker^[8] in sesame. The significant improvement in the number of pods plant⁻¹ in nipping at 45 DAS and 35 DAS might be due to initiation of higher number of branches plant⁻¹ which probably originated more flower buds that resulted in more pods. The results were in accordance with the findings of other workers^[2,6]. The increased yield components might be attributed to activation of lateral dormant buds by arresting the terminal growth through nipping which might have

facilitated the significant increase in the yield attributes. Similar findings were also reported by other worker^[10] in pigeon pea. The increase in yield attributes with nipping at 45 DAS and 35 DAS might also be due to enhanced branching and dispersion of carbohydrates towards auxiliary vegetative buds below nipped portion which might have helped in production of more number of pods plant⁻¹ and grain yield compared to no nipping.

The significantly higher seed yield (2567 kg ha⁻¹), straw yield (4037 kg ha⁻¹), biological yield (6604 kg ha⁻¹) were recorded with nipping practices at 45 DAS followed by nipping at 35 DAS (2331 kg ha⁻¹, 3581 kg ha⁻¹ and 5912, respectively) and the lowest (1945 kg ha⁻¹, 3377 kg ha⁻¹ and 5322 kg ha⁻¹, respectively) with no nipping practices. The higher yield under nipping treatments might be because of the development of auxiliary buds which is inhibited by indole acetic acid (IAA) produced in the apical portion. When the source of auxins is removed by nipping, the lateral branching gets activated which resulted in increased number of pods plant⁻¹ (Table 1) thereby increased seed yield. This was also explained by other workers

in sesame^[3,8]. Increase in yield with nipping as compared with no nipping clearly indicates that the energy which was previously used by plant to become taller was diverted towards grain formation. Grain yield depends on yield attributes *i.e.*,

Effect of varieties on yield attributes and yield

Varieties were also significantly influenced the yield attributes of chickpea (Table 1). Among varieties, *cv.* JG 12 produced significantly more number of pods (102 plant⁻¹) followed by *cvs.* RVG 201 (94.9 plant⁻¹), JG 36 (89.9 plant⁻¹) and the lowest in *cv.* RVG 202 (80.0 plant⁻¹). However, seed index (100 seeds weight) was observed in *cv.* RVG 201 (21.7g) followed by *cvs.* RVG 202 (20.7g), JG 36 (16.9g) and the significantly the lowest in *cv.* JG 12 (16.6g). Similarly, number of seeds pod⁻¹ did also not differ significantly among varieties. The differences in yield attributes observed among chickpea varieties might be due to their differences in growth habit and genetic yielding ability. Similar varietal difference in chickpea with respect to yield attributes was also reported by other workers^[1] in chickpea and cowpea.

Interaction effect on yield attributes and yield

Interaction between nipping and varieties significantly influenced yield attributes of chickpea. The interactional effect of nipping at 45 DAS with variety JG-12 recorded higher number of pods plant⁻¹ (112) over other treatment combinations. However, number of seeds pod⁻¹ did not differ significantly due to various interaction effects. Significantly, highest seed index (22.5g) was registered in the interaction of nipping at 35 DAS with variety RVG- 202 over other

number of branches (data not given) and number of pods plant⁻¹ that is why grain yield was higher with nipping at 45 DAS followed by nipping at 35 DAS as compared to no nipping. This was similar to the findings of other worker^[6].

The significantly higher seed yield (2506 kg ha⁻¹), straw yield (3862 kg ha⁻¹) and biological yield (6368 kg ha⁻¹) were recorded in *cv.* JG 12 followed by *cv.* RVG 201 (2315 kg ha⁻¹, 3738 kg ha⁻¹ and 6053, respectively), JG 36 (2217 kg ha⁻¹, 3557 kg ha⁻¹ and 5774, respectively) and significantly the lowest in *cv.* RVG 202 (2086 kg ha⁻¹, 3502 kg ha⁻¹ and 5588 kg ha⁻¹, respectively). This yield variation in respect of various varieties might be due to variation in pod bearing ability and vigorous growth. The higher yield could be attributed due to greater growth parameters *viz.*, plant height, number of primary and secondary branches more LAI (data not given) and cumulative effect of yield attributes (Table 1). The variety RVG-202 is a poor yielder because of its poor growth, yield attributes and canopy makeup. Similar results were also reported by other worker in pigeon pea^[10].

interactions. The increase in yield attributes noticed in the interaction of nipping with varieties was mainly due to accumulation of more photosynthates which were utilized for production of more number of productive secondary branches and more number of pods plant⁻¹ in chickpea. Similar results were also obtained by other worker in pigeon pea^[7,8,9].

Interaction between nipping and varieties was also significantly influenced

the grain yield, straw yield, biological yield. The interactional effect of nipping at 45 DAS with variety JG-12 recorded significantly highest grain yield (2789 kg ha^{-1}), straw yield (4339 kg ha^{-1}) and biological yield (7128 kg ha^{-1}) over other

Conclusion

The results of present study concluded that nipping at 45 DAS recorded significantly higher yield attributes and yields over nipping at 35 DAS and no nipping in chickpea varieties. Nipping in chickpea at 45 DAS increased the grain yield (kg ha^{-1}) by 32.0% and 10.1% over no nipping and nipping at 35 DAS, respectively. Among chickpea varieties, JG-12 was found significantly superior over other varieties in terms of yield attributes and seed yield. Variety JG-12 increased the grain yield (kg ha^{-1}) by 8.2 %, 13.0% and 20.1% over RVG-201, JG-36 and RVG-202, respectively. The

References

1. Aziz, M.A. and Rahman, M.M. (1994). Effect of date of sowing on yield and yield components of kabuli gram (*Cicer arietinum*). *Indian Journal of Agriculture Science*, 64(9): 624-626.
2. Baloch, M.S. and Zubair, M. (2010). Effect of nipping on growth and yield of chickpea. *The journal of animal & plant sciences*, 20(3): 208-210.
3. Bharathi, K., Panneerselvam, P. and Bhagya, H.P. (2014). Effect of clipping and plant growth regulator along with different kind of fertilizers on yield and yield parameters in sesame during monsoon period. *Indian Journal of Agricultural Research*, 48(3): 232-236.
4. Gnyandev, B., Kurdikeri, M.B. and Salimath, P.M. (2019). Effect of nipping and foliar spray of growth

treatment combinations. Increase in yield of varieties with nipping as compared to no nipping that the energy which was previously used by varieties to become taller was diverted towards grain formation due to nipping.

interactional effect between nipping at 45 DAS with cv. JG-12 was found superior over rest of the combination for yield attributes and yield. The combination of nipping at 45 DAS with cv. JG-12 increased the grain yield (kg ha^{-1}) by 4.5% to 56.6% over rest of the combinations of treatments. On the basis of results of current study, it may be recommended that the farmer can get maximum return by the adoption of nipping at 45 DAS. Among the varieties, cv. JG-12 is very suitable variety to get higher return. The nipping practices at 45 DAS with cv. JG-12 may be suitable to increase farmer's income.

regulators on plant growth, seed yield and quality in chickpea varieties, *Journal of Entomology and Zoology Studies*, 7(2): 318-321.

5. Gomez, K.A. and Gomez, A.A. (1984). *Statistical Procedures for Agriculture Research*. 2nd Edition. A Wiley Inter-Science Publication, New York (USA).
6. Khan, E.A., Hussain, I., Shreyar and Ahmad, H.B. (2018). Influence of nipping and foliar application of nutrients on growth and yield of chickpea in rainfed condition, *Legume Research*, 41(5): 740-744.
7. Khan, H., Gul, R. and Khan, N.U. (2017). Appraisal of interaction among nipping of chickpea (*Cicer arietinum* L.) genotypes and their correlated response for grain yield. *The Journal*

- of Animal & Plant Sciences*, 27(4): 1295-1302.
8. Kithan, L. and Singh, R. (2017). Effect of nipping, crop geometry and different levels of nitrogen on the growth and yield of sesame (*Sesamum indicum* L.). *Journal of Pharmacognosy and phytochemistry*, 6(4): 1089-1092.
 9. Math, G., Venkatesh, M.S., Balol, G. and Revanappa, S.B. (2021). Productivity and economics of pigeon pea genotypes as influenced by planting geometry, growth retardant, *Legume Research*, DOI: 10.18805/LR-4430.
 10. Teggelli, Ahmad, R.G., Z.B., Yusufali, N. and Patil, M.C. (2020). Influence of nipping technology on growth, yield and economics of pigeonpea cultivated under rainfed situation. *Journal of Phramacognosy and Phytochemistry*, 9(2): 886-888.